


## In the claims:

1. (Currently Amended) An N-[(piperazinyl)hetaryl]arylsulfonamide compound of the general formula I



in which

- R is oxygen, a group N-R<sup>3</sup> or a group CR<sup>3a</sup>R<sup>3b</sup>;
- Q is a bivalent, 6-membered heteroaromatic radical ~~which possesses 1 or 2 N atoms as ring members~~ selected from pyridindiyl and pyrimidindiyl, and which optionally carries one or two substituents R<sup>a</sup> which is/are selected, independently of each other, from halogen, CN, NO<sub>2</sub>, CO<sub>2</sub>R<sup>4</sup>, COR<sup>5</sup>, C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-haloalkyl, NH<sub>2</sub>, NHR<sup>6</sup>, NR<sup>6</sup>R<sup>7</sup> and C<sub>1</sub>-C<sub>4</sub>-haloalkoxy;
- Ar is phenyl or a 6-membered heteroaromatic radical ~~which possesses 1 or 2 N atoms as ring members~~ selected from pyridinyl and pyrimidinyl, and which optionally carries one or two substituents R<sup>b</sup>, which is/are selected from halogen, NO<sub>2</sub>, CN, CO<sub>2</sub>R<sup>4</sup>, COR<sup>5</sup>, NH<sub>2</sub>, NHR<sup>6</sup>, NR<sup>6</sup>R<sup>7</sup>, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-haloalkyl, C<sub>1</sub>-C<sub>6</sub>-alkoxy, C<sub>1</sub>-C<sub>6</sub>-haloalkoxy, C<sub>2</sub>-C<sub>6</sub>-alkenyl, C<sub>2</sub>-C<sub>6</sub>-alkynyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkoxy, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl-C<sub>1</sub>-C<sub>4</sub>-alkyl and C<sub>1</sub>-C<sub>4</sub>-haloalkyl, with it also being possible for two radicals R<sup>b</sup> which are bonded to adjacent C atoms of Ar to be together C<sub>3</sub>-C<sub>4</sub>-alkylene;
- n is 0, 1 or 2;
- R<sup>1</sup> is hydrogen, C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-haloalkyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl-C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-hydroxyalkyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy-C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>3</sub>-C<sub>4</sub>-alkenyl or C<sub>3</sub>-C<sub>4</sub>-alkynyl;
- R<sup>2</sup> is C<sub>1</sub>-C<sub>4</sub>-alkyl or, together with R<sup>1</sup>, is C<sub>2</sub>-C<sub>5</sub>-alkylene or, in the case of n = 2, the two radicals R<sup>2</sup> can together be C<sub>1</sub>-C<sub>4</sub>-alkylene;

R<sup>3</sup> is hydrogen or C<sub>1</sub>-C<sub>4</sub>-alkyl;

R<sup>3a</sup>, R<sup>3b</sup> are, independently of each other, hydrogen or C<sub>1</sub>-C<sub>4</sub>-alkyl;

R<sup>4</sup> is C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-haloalkyl, C<sub>2</sub>-C<sub>4</sub>-alkenyl C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl-C<sub>1</sub>-C<sub>4</sub>-alkyl, phenyl or benzyl; and

R<sup>5</sup> is hydrogen, C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-haloalkyl, C<sub>2</sub>-C<sub>4</sub>-alkenyl C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl-C<sub>1</sub>-C<sub>4</sub>-alkyl, phenyl or benzyl;

R<sup>6</sup>, R<sup>7</sup> are each independently selected from C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-haloalkyl or together with the nitrogen to which they are bound form a saturated 3-, 4-, 5- or 6-membered heterocycle, which additionally may comprise an oxygen atom or an additional nitrogen atom as a ring member and which may carry 1, 2, 3 or 4 C<sub>1</sub>-C<sub>4</sub> alkyl groups;

the N-oxides thereof and the physiologically tolerated acid addition salts of these compounds;

with the exception of the compounds: 4-methyl-N-[6-(4-methylpiperazin-1-yl)pyridin-3-yl]benzenesulfonamide and 4-chloro-N-[6-(4-methylpiperazin-1-yl)pyridin-3-yl]benzenesulfonamide.


2. (Original) The compound as claimed in claim 1, wherein R is N-R<sup>3</sup> with R<sup>3</sup> being H or C<sub>1</sub>-C<sub>4</sub>-alkyl.
3. (Currently Amended) The compound as claimed in claim 2, wherein

Q is a bivalent, 6-membered heteroaromatic radical ~~which possesses 1 or 2 N atoms as ring members~~ selected from pyridindiyl and pyrimidindiyl, and which optionally carries one or two substituents R<sup>a</sup> which is/are selected, independently of each other, from halogen, CN, NO<sub>2</sub>, CO<sub>2</sub>R<sup>4</sup>, COR<sup>5</sup>, C<sub>1</sub>-C<sub>4</sub>-alkyl and C<sub>1</sub>-C<sub>4</sub>-haloalkyl and

Ar is phenyl or a 6-membered heteroaromatic radical ~~which possesses 1 or 2 N atoms as ring members~~ selected from pyridinyl and pyrimidinyl, and which optionally carries one or two substituents R<sup>b</sup>, which is/are selected from halogen, NO<sub>2</sub>, CN, CO<sub>2</sub>R<sup>4</sup>, COR<sup>5</sup>, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>2</sub>-C<sub>6</sub>-alkenyl, C<sub>2</sub>-C<sub>6</sub>-alkynyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, C<sub>3</sub>-C<sub>6</sub>-


cycloalkyl-C<sub>1</sub>-C<sub>4</sub>-alkyl and C<sub>1</sub>-C<sub>4</sub>-haloalkyl, with it also being possible for two radicals R<sup>b</sup> which are bonded to adjacent C atoms of Ar to be together C<sub>3</sub>-C<sub>4</sub>-alkylene.

4. (Original) The compound as claimed in claim 1, in which the piperazine ring is bonded to the heteroaromatic radical Q in the para position in relation to the group R-SO<sub>2</sub>-Ar.
5. (Currently Amended) The compound as claimed in claim 1, in which Q is a radical of the formula




~~in which A<sub>1</sub>, A<sub>2</sub> and A<sub>3</sub> are, independently of each other, N or CH, one or two of the variables A<sub>1</sub>, A<sub>2</sub> and A<sub>3</sub> can also be C-R<sup>a</sup>, one of the variables A<sub>1</sub>, A<sub>2</sub> or A<sub>3</sub> is N, the remaining two variables being CH or C-R<sup>a</sup>, or A<sub>1</sub> and A<sub>3</sub> are N and A<sub>3</sub> is CH or C-R<sup>a</sup>, k = 0 or 1 and R<sup>a</sup> is selected from halogen, C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-haloalkyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy, NH<sub>2</sub>, NHR<sup>6</sup>, NR<sup>6</sup>R<sup>7</sup> and C<sub>1</sub>-C<sub>4</sub>-haloalkoxy, with A<sub>1</sub>, A<sub>2</sub> and A<sub>3</sub> not simultaneously being N or simultaneously being selected from CH and C-R<sup>a</sup> with the proviso that k is 0 if two of the variables A<sub>1</sub>, A<sub>2</sub> and A<sub>3</sub> are C-R<sup>a</sup>.~~

6. (Original) The compound as claimed in claim 5, in which A<sub>3</sub> is nitrogen, A<sub>2</sub> is CH and A<sub>1</sub> is N or CH and wherein the piperazine radical is located in the 2 position.
7. (Original) The compound as claimed in claim 6, in which Q is pyridin-2,5-diyl which carries the piperazine radical in the 2 position.
8. (Currently Amended) The compound as claimed in claim 6 5, in which Q is a radical of the formula



in which  $A_1$  and  $A_2$  are, independently of each other, N or CH  $A_1$  is N or CH and  $A_2$  is CH and  $R^a$  is selected from,  $C_1$ - $C_4$ -alkoxy,  $NH_2$ ,  $NHR^6$ ,  $NR^6R^7$  and  $C_1$ - $C_4$ -haloalkoxy.

9. (Currently Amended) The compound as claimed in claim 8, in which  ~~$A_1$  is N or CH and  $A_2$  is CH~~ and wherein the piperazine radical is located in the 2 position.
10. (Currently Amended) The compound as claimed in claim 1, in which the radical Ar carries a substituent  $R^b$  in the para position and, ~~where appropriate~~ optionally, a further substituent  $R^b$  in the meta position or in the ortho position, in each case based on the binding site of the sulfonamide group.
11. (Currently Amended) The compound as claimed in claim 1, in which Ar is phenyl or pyridyl, which radicals possess, ~~where appropriate~~, one or 2  $R^b$  substituents.
12. (Currently Amended) The compound as claimed in claim 1, in which  $R^1$  is ~~different from~~ not hydrogen and or methyl.
13. (Currently Amended) The compound as claimed in claim 1 of the general formula Ia




in which  $n$ ,  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^a$  and  $R^b$  have the meanings given in claim 1 and in which ~~either  $A_1$ ,  $A_2$  and  $A_3$  are, independently of each other, N or CH and one or two of the variables  $A_1$ ,  $A_2$  and  $A_3$  can also be  $C-R^a$ ; one of the variables  $A_1$ ,  $A_2$  or  $A_3$  is N, the remaining two variables being CH or  $C-R^a$ , or  $A_1$  and  $A_3$  are N and  $A_2$  is CH or  $C-R^a$ , with the proviso that  $k$  is 0 if two of the variables  $A_1$ ,  $A_2$  and  $A_3$  are  $C-R^a$ , with  $A_1$ ,  $A_2$  and  $A_3$  not simultaneously being N or simultaneously being selected from CH and  $C-R^a$ ;~~

X and Y are selected from CH,  $C-R^b$  and N, in which  $R^b$  is halogen, methyl, CN, difluoromethyl or trifluoromethyl, with X and Y not simultaneously being N or simultaneously being  $C-R^b$ , and


$k$  is 0 or 1.

14. (Currently Amended) The compound of the formula Ia as claimed in claim 13, in which  $k = 0$ , ~~with  $A_1$ ,  $A_2$  and  $A_3$  being, independently of each other, N or CH and  $A_1$ ,  $A_2$  and  $A_3$  not simultaneously being N or simultaneously being CH and one of the variables  $A_1$ ,  $A_2$  or  $A_3$  is N, the remaining two variables being CH or  $A_1$  and  $A_3$  are N and  $A_2$  is CH.~~
15. (Original) The compound of the formula Ia as claimed in claim 14, in which  $A_1$  is CH or N,  $A_2$  is CH and  $A_3$  is N.
16. (Original) The compound of the formula Ia as claimed in claim 13, in which  $k$  is 1,  $A_1$  is CH or N,  $A_2$  is CH and  $A_3$  is N, and  $R^a$  is selected from,  $C_1$ - $C_4$ -alkoxy,  $NH_2$ ,  $NHR^6$ ,  $NR^6R^7$  and  $C_1$ - $C_4$ -haloalkoxy and  $R^a$  is bound to the carbon atom adjacent to  $A_3$ .
17. (Previously Amended) The compound of the formula Ia as claimed in claim 13, in which  $n$  is 0 or 1 and, in the case of  $n = 1$ ,  $R^2$  is bonded to the C atom of the piperazine ring which is adjacent to the group  $R^1$ -N and is a methyl group having the S configuration.
18. (Currently Amended) The compound of the formula Ia as claimed in claim 13, in which the radical Ar carries a substituent  $R^b$  in the para position and, ~~where appropriate,~~ optionally a further substituent  $R^b$  in the meta position or in the ortho position, in each case based on the binding site of the sulfonamide group.
19. (Currently Amended) The compound of the formula Ia as claimed in claim 13, in which Ar is phenyl or pyridyl, which radicals possess, ~~where appropriate,~~ one or 2  $R^b$  substituents.
20. (Currently Amended) The compound of the formula Ia as claimed in claim 13, in which  $R^1$  is ~~different from~~ not hydrogen and or methyl.
21. (Previously Presented) The compound of the formula Ia as claimed in claim 13, of the general formula Ia.1



in which n, X, Y, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>a</sup> and R<sup>b</sup> have the meanings given in claim 13 and q is 0, 1 or 2.

22. (Previously Presented) The compound of the formula Ia as claimed in claim 13, of the general formula Ia.2



in which n, X, Y, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>a</sup> and R<sup>b</sup> have the meanings given in claim 13 and q is 0 or 1.

23. (Currently Amended) A pharmaceutical composition which comprises at least one N-[(piperazinyl)hetaryl]arylsulfonamide compound as claimed in claim 1 and/or at least one physiologically tolerated acid addition salt of I and/or an N-oxide of I, ~~where appropriate~~ optionally together with physiologically acceptable carriers and/or auxiliary substances.
24. (Canceled)
25. (Canceled)
26. (Canceled)
27. (Currently Amended) A method for treating a medical disorder susceptible to treatment with a dopamine D<sub>3</sub> receptor antagonist or a dopamine D<sub>3</sub> agonist selected from Parkinson's disease, schizophrenia, cognitive disturbances, depression, anxiety, addiction, kidney function disturbances, eating disturbances and epilepsy, said method comprising administering an effective amount of at least one compound of the formula I of claim 1 to a subject in need thereof.
28. (Canceled)